

On the Derived Functors of the Third Symmetric-Power Functor

Bernhard Köck and Ramesh Satkurunath

November 3, 2009

Abstract. We compute the derived functors of the third symmetric-power functor and their cross-effects for certain values. These calculations match predictions by the first named author and largely prove them in general.

Mathematics Subject Classification 2000. 13D25; 18G10; 18G30.

1 Introduction

Let R and S be rings. We recall that the construction of the left derived functors $L_k F : R\text{-}\mathbf{mod} \rightarrow S\text{-}\mathbf{mod}$ of any covariant right-exact functor $F : R\text{-}\mathbf{mod} \rightarrow S\text{-}\mathbf{mod}$ is achieved by applying three functors. The first functor constructs a projective resolution $P_\bullet$ of the R -module M that we wish to calculate the derived functor of. Then the functor F is applied to the resolution $P_\bullet$ giving the chain complex $F(P_\bullet)$. Lastly the k^{th} derived functor $L_k F$ is defined to be $H_k(F(P_\bullet))$, the k^{th} homology of the chain complex $F(P_\bullet)$. However for a given module M the projective resolution of M is unique only up to chain-homotopy equivalence, so this construction crucially depends on the fact that F preserves chain-homotopies. In general this fact does not hold when F is a nonlinear functor such as the l^{th} symmetric-power functor, Sym^l , or the l^{th} exterior-power functor, Λ^l . In the paper [DP] Dold and Puppe overcome this problem and define the derived functors of nonlinear functors by passing to the category of simplicial complexes using the Dold-Kan correspondence.

The Dold-Kan correspondence gives a pair of functors Γ and N that provide an equivalence between the category of bounded chain complexes and the category of simplicial complexes; under this correspondence chain homotopies correspond to simplicial homotopies. Furthermore in the simplicial world all functors preserve simplicial homotopy (not just linear functors).

Because of this the above definition of the derived functors of F becomes well defined for any functor when $F(P.)$ is replaced by the complex $NFT(P.)$.

Now let R be a Noetherian commutative ring, let I be an ideal in R which is locally generated by a regular sequence of length d , let V be a finitely generated projective R/I -module and let $P.(V)$ be an R -projective resolution of V . In [Kö] the first named author explicitly calculates the modules $H_k N \text{Sym}^l \Gamma P.(V)$ when $d = 1$ and l is any positive integer, and also when $d = 2$ and $l = 2$ (see Theorems 3.2 and 6.4 in [Kö]). As explained in Section 4 such calculations lie at the heart of a new approach to the seminal Adams-Riemann-Roch Theorem and hence to Grothendieck's Riemann-Roch theory.

In this paper we look at the case when $d = 2$ and $l = 3$. Throughout this paper we use the symbol G_k to denote the derived functor

$$\begin{aligned} G_k : \mathcal{P}_{R/I} &\rightarrow R\text{-}\mathbf{mod} \\ V &\mapsto H_k N \text{Sym}^3 \Gamma P.(V) \end{aligned}$$

where $\mathcal{P}_{R/I}$ denotes the category of finitely generated projective R/I -modules. Let L_1^3 denote the Schur functor indexed with Young diagram of shape $(2, 1)$ (see Definition 2.5). In Example 6.6 of [Kö] the first named author made the following prediction about the functor G_k

$$G_k(V) \cong \begin{cases} \text{Sym}^3(V) & k = 0 \\ L_1^3(V) \otimes I/I^2 & k = 1 \\ L_1^3(V) \otimes I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ D^3(V) \otimes \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \\ 0 & k \geq 5 \end{cases}$$

and for the case when $k = 2$ he suggests that there exists an exact sequence:

$$\begin{aligned} 0 \rightarrow D^2(V) \otimes V \otimes \Lambda^2(I/I^2) &\rightarrow H_2 N \text{Sym}^3 \Gamma(P.(V)) \\ &\rightarrow \Lambda^3(V) \otimes \text{Sym}^2(I/I^2) \rightarrow 0. \end{aligned}$$

Note that the prediction for $G_3(V)$ is given in slightly different terms but, as will be explained after Proposition 2.6, this formulation is equivalent.

For each k let F_k stand for the prediction made for G_k to be (note that F_2 is defined only on objects, not on morphisms). We will show that these predictions are true if V is the free module R/I of rank 1:

$$G_k(R/I) \cong F_k(R/I),$$

see Theorem 5.5. Moreover we will show that the following isomorphisms hold for the higher cross-effects (see Section 2 for the definition of cross-effect functors):

$$\begin{aligned} \mathrm{cr}_2(G_k)(R/I, R/I) &\cong \mathrm{cr}_2(F_k)(R/I, R/I) \\ \mathrm{cr}_3(G_k)(R/I, R/I, R/I) &\cong \mathrm{cr}_3(F_k)(R/I, R/I, R/I), \end{aligned}$$

see Proposition 2.8 and Theorems 4.1 and 3.2.

Theorem 1.5 of [Kö] implies that, if these isomorphisms commute with certain structural maps, then the predictions are true in general. To verify that the isomorphisms commute with the maps as required, we would need to not only complete some quite involved calculations, but we would also require a proper definition of the functor F_2 i.e. one which applies to objects and morphisms (rather than just to objects). Finding such a suitable candidate for F_2 remains an open problem.

2 Cross-effect Functors

In [EM] Eilenberg and Mac Lane introduced the theory of cross-effect functors. The theory of cross-effect functors are central to the calculations of this paper. In [Kö] the first named author proved a result (Theorem 1.5) which shows that two functors are isomorphic if they agree on certain data given by their cross-effect functors. In this section we set the scene by introducing some of the theory of cross-effect functors from [EM]. Then we introduce the aforementioned theorem from [Kö] (see Theorem 2.4) and exemplify it by showing that the Schur functor L_1^3 and the co-Schur functor $\tilde{L}_1^3$ are isomorphic. Furthermore we begin calculating the cross-effect functors of G_k and calculate $F_k(R/I)$, $\mathrm{cr}_2(F_k)(R/I, R/I)$ and $\mathrm{cr}_3(F_k)(R/I, R/I, R/I)$.

Let $\mathcal{P}$ be an additive category, let $\mathcal{M}$ be an abelian category, and let $F : \mathcal{P} \rightarrow \mathcal{M}$ be a functor with $F(0) = 0$.

Definition 2.1. Let $k \geq 0$. For any $V_1, \dots, V_k \in \mathcal{P}$ and $1 \leq i \leq k$, let

$$P_i : V_1 \oplus \dots \oplus V_k \rightarrow V_i \rightarrow V_1 \oplus \dots \oplus V_k$$

denote the i^{th} projection. The k -functor

$$\begin{aligned} \mathrm{cr}_k(F) : \quad \mathcal{P}^k &\rightarrow \mathcal{M} \\ (V_1, \dots, V_k) &\mapsto \mathrm{Im} \left(\sum_{j=1}^k \sum_{1 \leq i_1 \leq \dots \leq i_j \leq k} F(p_{i_1} + \dots + p_{i_j}) \right) \end{aligned}$$

is called the k^{th} cross-effect of F ; here for any $V_1, \dots, V_k, W_1, \dots, W_k \in \mathcal{P}$ and $f_1 \in \mathrm{Hom}_{\mathcal{P}}(V_1, W_1), \dots, f_k \in \mathrm{Hom}_{\mathcal{P}}(V_k, W_k)$, the map

$$\mathrm{cr}_k(F)(f_1, \dots, f_k) : \mathrm{cr}_k(F)(V_1, \dots, V_k) \rightarrow \mathrm{cr}_k(F)(W_1, \dots, W_k)$$

is induced by $f_1 \oplus \dots \oplus f_k \in \text{Hom}_{\mathcal{P}}(V_1 \oplus \dots \oplus V_k, W_1 \oplus \dots \oplus W_k)$. The functor F is said to be of *degree less than k* if $\text{cr}_k(F)$ is identically zero.

The cross-effect functors $\text{cr}_k(F), k \geq 0$, have the following properties. We obviously have $\text{cr}_0(F) \equiv 0$ and $\text{cr}_1(F) = F$. Furthermore, we have $\text{cr}_k(F)(V_1, \dots, V_k) = 0$ if $V_i = 0$ for any $i \in \{1, \dots, k\}$ (see Theorem 9.2 on p.79 in [EM]). The most important property of cross-effects is given in the following proposition.

Proposition 2.2. *For any $k, l \geq 1$ and $V_1, \dots, V_l \in \mathcal{P}$, we have a canonical isomorphism*

$$\text{cr}_k(F)(\dots, V_1 \oplus \dots \oplus V_l, \dots) \cong \bigoplus_{1 \leq j \leq l} \bigoplus_{1 \leq i_1 < \dots < i_j \leq k} \text{cr}_{k+j-1}(F)(\dots, V_{i_1}, \dots, V_{i_j}, \dots)$$

which is functorial in $V_1, \dots, V_l$. In particular, F is of degree $\leq k$, if and only if $\text{cr}_k(F)$ is a k -additive functor.

Proof. See Theorem 9.1, Lemma 9.8 and Lemma 9.9 in [EM]. $\square$

From Proposition 2.2 we see that

$$F(V_1 \oplus V_2) \cong F(V_1) \oplus F(V_2) \oplus \text{cr}_2(F)(V_1, V_2)$$

for all $V_1, V_2 \in \mathcal{P}$; i.e. $\text{cr}_2(F)$ measures the deviation from linearity of the functor F . This isomorphism can also be used to define $\text{cr}_2(F)$ (see section 3 in [JM]). Similarly, the higher cross-effects can be defined inductively by the isomorphism

$$\begin{aligned} \text{cr}_k(F)(V_1, \dots, V_{k-1}, V_k \oplus V'_k) &\cong \text{cr}_k(F)(V_1, \dots, V_{k-1}, V_k) \\ &\quad \oplus \text{cr}_k(F)(V_1, \dots, V_{k-1}, V'_k) \\ &\quad \oplus \text{cr}_{k+1}(F)(V_1, \dots, V_{k-1}, V_k, V'_k) \end{aligned}$$

(see section 7 in [JM]). When actually calculating cross-effect functors it is this definition that we will use.

Definition 2.3. Let $l \geq k \geq 1$, $V_1, \dots, V_k \in \mathcal{P}$, and let $\epsilon = (\epsilon_1, \dots, \epsilon_k) \in \{1, \dots, l\}^k$ with $|\epsilon| := \sum_{i=1}^k \epsilon_i = l$. The composition

$$\begin{aligned} \Delta_\epsilon : \text{cr}_k(F)(V_1, \dots, V_k) &\xrightarrow{\text{cr}_k(F)(\Delta, \dots, \Delta)} \text{cr}_k(F)(V_1^{\epsilon_1}, \dots, V_k^{\epsilon_k}) \\ &\xrightarrow{\pi} \text{cr}_l(F)(V_1, \dots, V_1, \dots, V_k, \dots, V_k) \end{aligned}$$

of the map $\text{cr}_k(F)(\Delta, \dots, \Delta)$ (induced by the diagonal maps $\Delta : V_i \rightarrow V_i^{\epsilon_i}, i = 1, \dots, k$) with the canonical projection π (according to Proposition 2.2) is called the *diagonal map associated with ϵ* . The analogous composition

$$\begin{aligned} +_\epsilon : \text{cr}_l(F)(V_1, \dots, V_1, \dots, V_k, \dots, V_k) &\hookrightarrow \text{cr}_k(F)(V_1^{\epsilon_1}, \dots, V_k^{\epsilon_k}) \\ &\xrightarrow{\text{cr}_k(F)(+, \dots, +)} \text{cr}_k(F)(V_1, \dots, V_k) \end{aligned}$$

is called *plus map associated with ϵ* .

The maps Δ_ϵ and $+_\epsilon$ form natural transformations between the functors $\text{cr}_k(F)$ and $\text{cr}_l(F) \circ (\Delta_{\epsilon_1}, \dots, \Delta_{\epsilon_k})$ from $\mathcal{P}^k$ to $\mathcal{M}$. One easily sees that the map Δ_ϵ can be decomposed into a composition of maps Δ_δ with $\delta \in \{1, 2\}^j$ such that $|\delta| = j + 1$ and $j \in \{k, \dots, l - 1\}$. The same holds for $+_\epsilon$.

Theorem 2.4. *Let A be a ring, $\mathcal{M}$ an abelian category, $d \in \mathbb{N}_+$, and*

$$F, G : (\text{f.g. projective } A\text{-modules}) \rightarrow \mathcal{M}$$

be two functors of degree $\leq d$ with $F(0) = 0 = G(0)$. Suppose that there exist isomorphisms

$$\alpha_i(A, \dots, A) : \text{cr}_i(F)(A, \dots, A) \xrightarrow{\sim} \text{cr}_i(G)(A, \dots, A), \quad i = 1, \dots, d,$$

which are compatible with the action of A in each component and which make the following diagrams commute for $i \in \{1, \dots, d - 1\}$ and $\epsilon \in \{1, 2\}^i$ with $|\epsilon| = i + 1$:

$$\begin{array}{ccc} \text{cr}_i(F)(A, \dots, A) & \longrightarrow & \text{cr}_i(G)(A, \dots, A) \\ \downarrow \Delta_\epsilon & & \downarrow \Delta_\epsilon \\ \text{cr}_{i+1}(F)(A, \dots, A) & \longrightarrow & \text{cr}_{i+1}(G)(A, \dots, A) \\ \\ \text{cr}_{i+1}(F)(A, \dots, A) & \longrightarrow & \text{cr}_{i+1}(G)(A, \dots, A) \\ \downarrow +_\epsilon & & \downarrow +_\epsilon \\ \text{cr}_i(F)(A, \dots, A) & \longrightarrow & \text{cr}_i(G)(A, \dots, A). \end{array}$$

Then the two functors F and G are isomorphic.

Proof. See Theorem 1.5 of [Kö]. □

Definition 2.5. Let A be a commutative ring. For any finitely generated A -module V we define $L_1^3(V)$ by the following exact sequences and we call L_1^3 the Schur functor indexed by the Young diagram of $(2, 1)$.

$$\begin{array}{ccccccc}
\Lambda^3(V) & \longrightarrow & \Lambda^2(V) \otimes V & \longrightarrow & V \otimes \text{Sym}^2(V) & \longrightarrow & \text{Sym}^3(V) \\
& & \searrow & & \nearrow & & \\
& & L_1^3(V) & & & & \\
& \nearrow & & \searrow & & & \\
0 & & & & & & 0
\end{array}$$

And we define the co-Schur functor $\tilde{L}_1^3$ by the following exact sequence.

$$\begin{array}{ccccccc}
D^3(V) & \longrightarrow & D^2(V) \otimes V & \longrightarrow & V \otimes \Lambda^2(V) & \longrightarrow & \Lambda^3(V) \\
& & \searrow & & \nearrow & & \\
& & \tilde{L}_1^3(V) & & & & \\
& \nearrow & & \searrow & & & \\
0 & & & & & & 0
\end{array}$$

Here, the horizontal sequences are the well-known exact Koszul and co-Koszul complexes, see Definition 5.1.

We now give an example of how Theorem 2.4 can be used, by proving that the Schur functor L_1^3 and the co-Schur functor $\tilde{L}_1^3$ are isomorphic.

Proposition 2.6.

$$\text{cr}_k(L_1^3)(A, \dots, A) \cong \text{cr}_k(\tilde{L}_1^3)(A, \dots, A) \cong \begin{cases} 0 & k = 1 \\ A \oplus A & k = 2 \\ A \oplus A & k = 3 \\ 0 & k \geq 4. \end{cases}$$

Moreover the two functors L_1^3 and $\tilde{L}_1^3$ are isomorphic.

Note this justifies the reformulation of the predictions for G_k as seen in the introduction, as the original prediction for G_3 in [Kö] was written as $G_3(V) \cong \tilde{L}_1^3(V) \otimes I/I^2 \otimes \Lambda^2(I/I^2)$.

Proof. Let V and W be finitely generated projective A -modules. Definition 2.5 tells us the co-Schur functor $\tilde{L}_1^3(V)$ is given by the short exact sequence

$0 \rightarrow \tilde{L}_1^3(V) \rightarrow V \otimes \Lambda^2(V) \rightarrow \Lambda^3(V) \rightarrow 0$. In particular this tells us that $\text{cr}_1(\tilde{L}_1^3)(A) = 0$, similarly we see that $L_1^3(A) = 0$.

Next we want to compute $\text{cr}_2(\tilde{L}_1^3)(A, A)$ and $\text{cr}_2(L_1^3)(A, A)$. We have the short exact sequence $0 \rightarrow L_1^3(V \oplus W) \rightarrow (V \oplus W) \otimes \Lambda^2(V \oplus W) \rightarrow \Lambda^3(V \oplus W) \rightarrow 0$. Using the canonical decomposition for Λ^n we get the following short exact sequence:

$$\begin{aligned} 0 \rightarrow \text{cr}_2(\tilde{L}_1^3)(V, W) \rightarrow \\ V \otimes (V \otimes W) \oplus V \otimes \Lambda^2(W) \oplus W \otimes \Lambda^2(V) \oplus W \otimes (V \otimes W) \\ \rightarrow \Lambda^2(V) \otimes W \oplus V \otimes \Lambda^2(W) \rightarrow 0. \end{aligned}$$

Therefore $\text{cr}_2(\tilde{L}_1^3)(V, W) \cong V^{\otimes 2} \otimes W \oplus V \otimes W^{\otimes 2}$. Similarly we find that $\text{cr}_2(L_1^3)(V, W) \cong V^{\otimes 2} \otimes W \oplus V \otimes W^{\otimes 2}$, so $\text{cr}_2(\tilde{L}_1^3)$ and $\text{cr}_2(L_1^3)$ are isomorphic as bi-functors. Because of the way that higher cross-effects are calculated from lower cross-effects we see that, for $k \geq 2$, the k -functors $\text{cr}_k(\tilde{L}_1^3)$ and $\text{cr}_k(L_1^3)$ will be isomorphic and the corresponding diagrams of $+$ and Δ maps will commute. We easily verify that $\text{cr}_2(L_1^3)(A, A) \cong \text{cr}_2(\tilde{L}_1^3)(A, A)$ and $\text{cr}_3(L_1^3)(A, A, A) \cong \text{cr}_3(\tilde{L}_1^3)(A, A, A)$ are free of rank 2 over A , as stated.

Since $\text{cr}_1(\tilde{L}_1^3)(A) \cong \text{cr}_1(L_1^3)(A) \cong 0$ the corresponding $+$ and Δ maps involving $\text{cr}_1(\tilde{L}_1^3)(A)$ and $\text{cr}_1(L_1^3)(A)$ will just be zero maps, and hence will commute with the isomorphism. So Theorem 2.4 implies that $\tilde{L}_1^3$ is isomorphic to L_1^3 . $\square$

The following lemma begins the work calculating the cross-effects of G_k that we will complete in the following chapters. Let A and A' be simplicial R -modules; the reader should be aware that in the following lemma and its proof that $A \otimes A'$ means the simplicial module whose n^{th} place is $A_n \otimes A'_n$.

Lemma 2.7.

$$\begin{aligned} \text{cr}_2(G_k)(V, W) &\cong H_k N(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W)) \\ &\quad \oplus H_k N(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W)) \\ \text{cr}_3(G_k)(V, W, X) &\cong H_k N(\Gamma P.(V) \otimes \Gamma P.(W) \otimes \Gamma P.(X)). \end{aligned}$$

Proof. First we calculate $G_k(V \oplus W)$ for R -modules V, W to give us an expression for $\text{cr}_2(G_k)(V, W)$. To do this we use the fact that P, Γ, N and H_k are linear functors and also the canonical decomposition $\text{Sym}^3(V \oplus W) \cong$

$$\bigoplus_{l=0}^3 \text{Sym}^{3-l}(V) \otimes \text{Sym}^l(W).$$

$$\begin{aligned} G_k(V \oplus W) &= H_k N \text{Sym}^3 \Gamma P.(V \oplus W) \\ &\cong H_k N \left(\text{Sym}^3 \Gamma P.(V) \oplus \text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W) \right. \\ &\quad \left. \oplus \Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W) \oplus \text{Sym}^3 \Gamma P.(W) \right) \\ &\cong G_k(V) \oplus H_k N \left(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W) \right) \\ &\quad \oplus H_k N \left(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W) \right) \oplus G_k(W). \end{aligned}$$

And hence we get the desired expression for $\text{cr}_2(G_k)(V, W)$.

We now complete the proof by using our expression for $\text{cr}_2(G_k)(V, W)$, to get an expression for $\text{cr}_3(G_k)(V, W, X)$. We start this by calculating $\text{cr}_2(G_k)(V, W \oplus X)$ for R -modules V, W, X . To simplify our calculation we split up our expression for $\text{cr}_2(G_k)(V, W \oplus X)$ into $H_k N(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W \oplus X))$ and $H_k N(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W \oplus X))$, calculate each part separately, then add them together afterwards.

$$\begin{aligned} H_k N \left(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W \oplus X) \right) \\ \cong H_k N \left(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W) \right) \\ \oplus H_k N \left(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(X) \right). \end{aligned}$$

$$\begin{aligned} H_k N \left(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W \oplus X) \right) \\ \cong H_k N \left(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W) \right) \\ \oplus H_k N \left(\Gamma P.(V) \otimes (\Gamma P.(W) \otimes \Gamma P.(X)) \right) \\ \oplus H_k N \left(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(X) \right). \end{aligned}$$

This gives us the following expression for $\text{cr}_2(G_k)(V, W \oplus X)$:

$$\begin{aligned} \text{cr}_2(G_k)(V, W \oplus X) &\cong \text{cr}_2(G_k)(V, W) \oplus \text{cr}_2(G_k)(V, X) \\ &\quad \oplus H_k N \left(\Gamma P.(V) \otimes (\Gamma P.(W) \otimes \Gamma P.(X)) \right). \end{aligned}$$

Therefore $\text{cr}_3(G_k)(V, W, X) \cong H_k N(\Gamma P.(V) \otimes \Gamma P.(W) \otimes \Gamma P.(X))$ as desired. $\square$

We finally calculate the cross-effects of the predictions evaluated at the free R/I -module of rank 1.

Proposition 2.8. *We have the following R/I -module isomorphisms:*

$$F_k(R/I) \cong \begin{cases} R/I & k = 0 \\ 0 & k = 1 \\ \Lambda^2(I/I^2) & k = 2 \\ 0 & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \end{cases}$$

$$\begin{aligned} & \text{cr}_2(F_k)(R/I, R/I) \\ & \cong \begin{cases} R/I \oplus R/I & k = 0 \\ I/I^2 \oplus I/I^2 & k = 1 \\ \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) & k = 2 \\ I/I^2 \otimes \Lambda^2(I/I^2) \oplus I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} \oplus \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \end{cases} \end{aligned}$$

$$\begin{aligned} & \text{cr}_3(F_k)(R/I, R/I, R/I) \\ & \cong \begin{cases} R/I & k = 0 \\ I/I^2 \oplus I/I^2 & k = 1 \\ \Lambda^2(I/I^2) \oplus (I/I^2)^{\otimes 2} \oplus \Lambda^2(I/I^2) \oplus \text{Sym}^2(I/I^2) & k = 2 \\ I/I^2 \otimes \Lambda^2(I/I^2) \oplus I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} & k = 4. \end{cases} \end{aligned}$$

Proof. These results follow from a simple application of Proposition 2.6, and the canonical decompositions for the symmetric-, exterior-, and divided-power functors. $\square$

3 The Iterated Eilenberg-Zilber Theorem and Calculating $\text{cr}_3(G_k)(R/I, R/I, R/I)$

Let A^1 and A^2 be simplicial R -modules. The Eilenberg-Zilber Theorem (see §28 of [M], specifically Corollary 29.6 on p132) tells us that $N\Delta(A^1 \otimes A^2)$ is chain homotopic to $\text{Tot}(NA^1 \otimes NA^2)$ (where $\Delta(A^1 \otimes A^2)$ denotes the diagonal of the bisimplicial complex $A^1 \otimes A^2$). Let C^1, C'^1, C^2, C'^2 be chain

complexes of R -modules. If C^1 is chain homotopic to C'^1 and C^2 is chain homotopic to C'^2 then $\text{Tot}(C^1 \otimes C^2) \cong \text{Tot}(C'^1 \otimes C'^2)$; because of this the Eilenberg-Zilber Theorem can be iterated to give us the following theorem.

Theorem 3.1. The Iterated Eilenberg-Zilber Theorem. *Let $n \in \mathbb{N}$ with $n \geq 2$ and let $A^1, \dots, A^n$ be simplicial complexes. Then the complexes $N\Delta(A^1 \otimes \dots \otimes A^n)$ and $\text{Tot}(NA^1 \otimes \dots \otimes NA^n)$ are chain homotopic and (consequently) they are quasi-isomorphic:*

$$H_k(N\Delta(A^1 \otimes \dots \otimes A^n)) \cong H_k(\text{Tot}(NA^1 \otimes \dots \otimes NA^n)).$$

The following theorem, together with the canonical decomposition for exterior powers, shows that the third cross-effect functor for the derived functors of Sym^3 evaluated on $(R/I, R/I, R/I)$ matches the predictions (see Proposition 2.8).

Theorem 3.2.

$$\text{cr}_3(G_k)(R/I, R/I, R/I) \cong \Lambda^k(I/I^2 \oplus I/I^2)$$

Proof. From Lemma 2.7 we know that

$$\text{cr}_3(G_k)(V, W, X) \cong H_k N(\Gamma P.(V) \otimes \Gamma P.(W) \otimes \Gamma P.(X)).$$

Here $\Gamma P.(V) \otimes \Gamma P.(W) \otimes \Gamma P.(X)$ stands for the simplicial complex whose k^{th} place is $\Gamma P_k(V) \otimes \Gamma P_k(W) \otimes \Gamma P_k(X)$. But we may consider this to be the diagonal of the tri-simplicial complex whose $(k, l, m)^{\text{th}}$ place is $\Gamma P_k(V) \otimes \Gamma P_l(W) \otimes \Gamma P_m(X)$, and by doing so we write

$$\text{cr}_3(G_k)(V, W, X) \cong H_k N\Delta(\Gamma P.(V) \otimes \Gamma P.(W) \otimes \Gamma P.(X)).$$

The Iterated Eilenberg-Zilber Theorem tells us that

$$H_k N\Delta(\Gamma P.(R/I) \otimes \Gamma P.(R/I) \otimes \Gamma P.(R/I)) \cong H_k \text{Tot}(P.(R/I) \otimes P.(R/I) \otimes P.(R/I)).$$

Theorem 5.1 of [Kö] tells us that

$$H_k \text{Tot}(P.(R/I) \otimes P.(R/I) \otimes P.(R/I)) \cong \Lambda^k(I/I^2 \oplus I/I^2),$$

as desired. □

4 The Derived Functors of Sym^2 and Calculating $\text{cr}_2(G_k)(R/I, R/I)$

The following theorem shows that the second cross-effect functor of the derived functors of Sym^3 evaluated on $(R/I, R/I)$ matches the predictions (see Proposition 2.8). Essential to the following proof is the information that has already been calculated about $H_k N \text{Sym}^2 \Gamma P.$ in [Kö], and the use of the Hypertor functor to exploit this.

Theorem 4.1.

$$\text{cr}_2(G_k)(R/I, R/I) \cong \begin{cases} R/I \oplus R/I & k = 0 \\ I/I^2 \oplus I/I^2 & k = 1 \\ \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) & k = 2 \\ I/I^2 \otimes \Lambda^2(I/I^2) \oplus I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} \oplus \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \\ 0 & k \geq 5. \end{cases}$$

Proof. Lemma 2.7 tells us that

$$\begin{aligned} \text{cr}_2(G_k)(V, W) &\cong H_k N(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W)) \\ &\quad \oplus H_k N(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W)). \end{aligned}$$

Here $\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W)$ and $\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W)$ stand for simplicial modules whose k^{th} place is $\text{Sym}^2 \Gamma P_k(V) \otimes \Gamma P_k(W)$ and $\Gamma P_k(V) \otimes \text{Sym}^2 \Gamma P_k(W)$ respectively; however we may consider them to be the diagonal of the bi-simplicial modules whose $(k, l)^{\text{th}}$ place is $\text{Sym}^2 \Gamma P_k(V) \otimes \Gamma P_l(W)$ and $\Gamma P_k(V) \otimes \text{Sym}^2 \Gamma P_l(W)$ respectively, therefore we can write

$$\begin{aligned} \text{cr}_2(G_k)(V, W) &\cong H_k N \Delta(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W)) \\ &\quad \oplus H_k N \Delta(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W)). \end{aligned}$$

We now start to calculate $H_k N \Delta(\text{Sym}^2 \Gamma P.(V) \otimes \Gamma P.(W))$, later we will calculate $H_k N \Delta(\Gamma P.(V) \otimes \text{Sym}^2 \Gamma P.(W))$ and then add the two together to get an expression for $\text{cr}_2(G_k)(V, W)$.

Using the Eilenberg-Zilber Theorem (Theorem 3.1) we see that

$$\begin{aligned} H_k N \Delta(\text{Sym}^2(\Gamma P.(V) \otimes \Gamma P.(W))) &\cong H_k \text{Tot}(N \text{Sym}^2 \Gamma P.(V) \otimes N \Gamma P.(W)) \\ &\cong H_k \text{Tot}(N \text{Sym}^2 \Gamma P.(V) \otimes P.(W)). \end{aligned}$$

So we want to calculate $H_k \text{Tot}(N \text{Sym}^2 \Gamma P.(V) \otimes P.(W))$, but this is just the definition of the hypertor $\mathbf{Tor}_i^R(N \text{Sym}^2 \Gamma P.(V), W)$. Application 5.7.8 of [W] gives us a spectral sequence to calculate hypertor

$${}^{II}E_{pq}^2 = \text{Tor}_p(H_q(A), B) \Rightarrow \mathbf{Tor}_{p+q}^R(A_*, B).$$

Theorem 6.4 of [Kö] tells us that

$$H_k N \text{Sym}^2 \Gamma(P.(V)) \cong \begin{cases} \text{Sym}^2(V) & k = 0 \\ \Lambda^2(V) \otimes I/I^2 & k = 1 \\ D^2(V) \otimes \Lambda^2(I/I^2) & k = 2 \\ 0 & k \geq 3. \end{cases}$$

Now $\text{Sym}^2(R/I) \cong R/I$, $\Lambda^2(R/I) \cong 0$ and $D^2(R/I) \cong R/I$. Hence the terms in the 2nd sheet of our spectral sequence are $\text{Tor}_p(R/I, R/I)$ in the zeroth column, $\text{Tor}_p(0, R/I) = 0$ in the first column, $\text{Tor}_p(\Lambda(I/I^2), R/I)$ in the second column and 0 everywhere else. From Example 5.2 of [Kö] (also see Theorem 5.1 of [Kö]) we know that $\text{Tor}_k(R/I, R/I) \cong \Lambda^k(I/I^2)$ and hence have $\text{Tor}_k(V, W) \cong V \otimes W \otimes \Lambda^k(I/I^2)$. So the 2nd sheet of the spectral sequence looks like this:

$$\begin{array}{ccccccccc} & \ddots & & \vdots & & \vdots & & \vdots & & \vdots \\ \cdots & 0 & & 0 & & 0 & & 0 & & 0 \\ \cdots & 0 & & \Lambda^2(I/I^2)^{\otimes 2} & & 0 & & \Lambda^2(I/I^2) & & \\ \cdots & 0 & & I/I^2 \otimes \Lambda^2(I/I^2) & & 0 & & I/I^2 & & \\ \cdots & 0 & & \Lambda^2(I/I^2) & & 0 & & R/I. & & \end{array}$$

The differentials on this level of the spectral sequence are -2 in the p -direction and $+1$ in the q -direction, so each differential either comes from or goes to a zero module. Hence the differentials are all zero maps i.e. the spectral sequence has already converged on the second level.

Therefore

$$H_k N\Delta(\operatorname{Sym}^2 \Gamma P.(R/I) \otimes \Gamma P.(R/I)) \cong \begin{cases} R/I & k = 0 \\ I/I^2 & k = 1 \\ \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) & k = 2 \\ I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \\ 0 & k \geq 5. \end{cases}$$

By symmetry

$$H_k N\Delta(\Gamma P.(R/I) \otimes \operatorname{Sym}^2 \Gamma P.(R/I)) \cong H_k N\Delta(\operatorname{Sym}^2 \Gamma P.(R/I) \otimes \Gamma P.(R/I)).$$

Hence

$$\operatorname{cr}_2(G_k)(R/I, R/I) \cong \begin{cases} R/I \oplus R/I & k = 0 \\ I/I^2 \oplus I/I^2 & k = 1 \\ \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) \oplus \Lambda^2(I/I^2) & k = 2 \\ I/I^2 \otimes \Lambda^2(I/I^2) \oplus I/I^2 \otimes \Lambda^2(I/I^2) & k = 3 \\ \Lambda^2(I/I^2)^{\otimes 2} \oplus \Lambda^2(I/I^2)^{\otimes 2} & k = 4 \\ 0 & k \geq 5. \end{cases}$$

□

5 The Cauchy Decomposition of $\operatorname{Sym}^3(P \otimes Q)$ and Calculating $G_k(R/I)$

In this final section we complete our calculations of G_k , but before we proceed we introduce the Cauchy decomposition given in chapter III of [ABW] as it applies to the third symmetric power, and remind the reader of Koszul and co-Koszul complexes.

Let R be a ring. Let P and Q be finitely generated projective R -modules. We now summarize the Cauchy decomposition of $\operatorname{Sym}^3(P \otimes Q)$. This decomposition plays a central role in Theorem 5.5, our calculation of $G_k(R/I)$.

A three step filtration

$$0 \subset M_{(3)}(\operatorname{Sym}^3(P \otimes Q)) \subset M_{(2,1)}(\operatorname{Sym}^3(P \otimes Q)) \subset \operatorname{Sym}^3(P \otimes Q)$$

is put on $\text{Sym}^3(P \otimes Q)$. The $M_{(3)}(\text{Sym}^3(P \otimes Q))$ part is defined to be the image of the determinant map

$$\Lambda^3 P \otimes \Lambda^3 Q \rightarrow \text{Sym}^3(P \otimes Q)$$

$$p_1 \wedge p_2 \wedge p_3 \otimes q_1 \wedge q_2 \wedge q_3 \mapsto \begin{vmatrix} p_1 \otimes q_1 & p_1 \otimes q_2 & p_1 \otimes q_3 \\ p_2 \otimes q_1 & p_2 \otimes q_2 & p_2 \otimes q_3 \\ p_3 \otimes q_1 & p_3 \otimes q_2 & p_3 \otimes q_3 \end{vmatrix}$$

(note this is simply isomorphic to $\Lambda^3 P \otimes \Lambda^3 Q$). The $M_{(2,1)}(\text{Sym}^3(P \otimes Q))$ part is defined to be equal to the previous part $M_{(3)}(\text{Sym}^3(P \otimes Q))$ part plus the image of the following homomorphism:

$$\Lambda^2 P \otimes P \otimes \Lambda^2 Q \otimes Q \rightarrow \text{Sym}^3(P \otimes Q)$$

$$p_1 \wedge p_2 \otimes p_3 \otimes q_1 \wedge q_2 \otimes q_3 \mapsto \begin{vmatrix} p_1 \otimes q_1 & p_1 \otimes q_2 \\ p_2 \otimes q_1 & p_2 \otimes q_2 \end{vmatrix} (p_3 \otimes q_3).$$

The quotients of this filtration are given by the short exact sequences

$$0 \rightarrow \Lambda^3 P \otimes \Lambda^3 Q \rightarrow M_{(2,1)}(\text{Sym}^3(P \otimes Q)) \rightarrow L_1^3 P \otimes L_1^3 Q \rightarrow 0$$

and

$$0 \rightarrow M_{(2,1)}(\text{Sym}^3(P \otimes Q)) \rightarrow \text{Sym}^3(P \otimes Q) \rightarrow \text{Sym}^3 P \otimes \text{Sym}^3 Q \rightarrow 0.$$

We now properly introduce the definition of Koszul complexes, which we use for projective resolutions.

Definition 5.1. Let $f : P \rightarrow Q$ be a homomorphism between two finitely generated projective R -modules, and $n \in \mathbb{N}$. Let $\text{Kos}^n(f)$ be the *Koszul complex*

$$0 \rightarrow \Lambda^n P \xrightarrow{d_{n-1}} \Lambda^{n-1} P \otimes Q \xrightarrow{d_{n-2}} \dots \xrightarrow{d_1} P \otimes \text{Sym}^{n-1} Q \xrightarrow{d_0} \text{Sym}^n Q \rightarrow 0$$

where, for $k \in \{0, 1, \dots, n-1\}$, the differential

$$d_k : \Lambda^{k+1} P \otimes \text{Sym}^{n-k-1} Q \rightarrow \Lambda^k P \otimes \text{Sym}^{n-k} Q$$

acts by

$$p_1 \wedge \dots \wedge p_{k+1} \otimes q_{k+2} \dots q_n \mapsto \sum_{i=1}^{k+1} (-1)^{k+1-i} p_1 \wedge \dots \wedge \hat{p}_i \wedge \dots \wedge p_{k+1} \otimes f(p_i) q_{k+2} \dots q_n.$$

Now let $f^* : Q^* \rightarrow P^*$ denote the dual map, then the part of the Koszul complex $\text{Kos}^n(f^*)$ in the k^{th} degree is $\Lambda^k Q^* \otimes \text{Sym}^{n-k} P^*$. The dual of this chain complex is a co-chain complex with the part in the k^{th} degree being $(\Lambda^k Q^* \otimes \text{Sym}^{n-k} P^*)^* \cong \Lambda^k Q \otimes D^{n-k} P$, i.e.

$$0 \leftarrow \Lambda^n Q \leftarrow \Lambda^{n-1} Q \otimes P \leftarrow \dots \leftarrow Q \otimes D^{n-1} P \leftarrow D^n P \leftarrow 0.$$

We call this the *co-Koszul complex* and denote it by $\tilde{\text{Kos}}^n(f)$.

Remark 5.2. It is well known that the complexes $\text{Kos}(f)$ and $\tilde{\text{Kos}}(f)$ are exact if f is an isomorphism.

The following two propositions will be useful in the proof of Theorem 5.5.

Proposition 5.3. *Let $f : P \rightarrow Q$ be a homomorphism between two finitely generated projective R -modules. If we consider $P \rightarrow Q$ to be a chain complex concentrated in degrees 1 and 0 then we have quasi-isomorphisms*

$$\text{Kos}^n(f) \cong N \text{Sym}^n \Gamma(P \rightarrow Q)$$

and

$$\tilde{\text{Kos}}^n(f) \cong N \Lambda^n \Gamma(P \rightarrow Q),$$

where Γ and N are the functors of the Dold-Kan correspondence.

Proof. See Proposition 2.4 and Remark 3.6 of [Kö]. □

Proposition 5.4. *Let $A.$ and $B.$ be bounded complexes of finitely generated projective R -modules. Then we have a spectral sequence*

$${}^I E_{p,q}^2 = \bigoplus_{q=q'+q''} \text{Tor}_p(H_{q'}(A.), H_{q''}(B.)) \Rightarrow \mathbf{Tor}_{p+q}(A., B.)$$

where $\mathbf{Tor}_n(A., B.)$ is defined as $H_n \text{Tot}(A. \otimes B.)$. In particular if $A.'$ and $B.'$ are further complexes as above and $A \rightarrow A'$ and $B \rightarrow B'$ are quasi-isomorphisms then the induced morphism

$$\text{Tot}(A. \otimes B.) \rightarrow \text{Tot}(A.' \otimes B.)$$

is a quasi-isomorphism as well.

Proof. See Application 5.7.8 of [W]. □

The following theorem shows that if I is globally generated by a regular sequence then the derived functors of Sym^3 evaluated on R/I match the predictions.

Theorem 5.5. *If I is generated by a regular sequence of length 2 then the module $G_k(R/I)$ is a free R/I -module of rank 1, for $k = 0, 2$ or 4 and otherwise of rank 0.*

Proof. Let f, g be a regular sequence in R , and let I be generated by it. Also we let $K.$ denote the complex $\dots \rightarrow 0 \rightarrow R \xrightarrow{f} R$ and $L.$ denote the Koszul complex $\dots \rightarrow 0 \rightarrow R \xrightarrow{g} R$. We use the complex

$$\text{Kos}^2(R \oplus R \xrightarrow{(f,g)} R) \cong \text{Tot}(K. \otimes L.)$$

as a resolution of R/I .

We see that:

$$\begin{aligned} G_k(R/I) &:= H_k N \operatorname{Sym}^3 \Gamma \operatorname{Tot}(K. \otimes L.) \\ &\cong H_k N \operatorname{Sym}^3 \Gamma \operatorname{Tot}(N\Gamma K. \otimes N\Gamma L.). \end{aligned}$$

Theorem 3.1 tells us that $\operatorname{Tot}(N\Gamma K. \otimes N\Gamma L.)$ is chain homotopic to $N\Delta(\Gamma K. \otimes \Gamma L.)$. Applying Γ turns the notion of chain homotopy into simplicial homotopy, all functors preserve homotopy in the simplicial world and N changes the notion of simplicial homotopy into the notion of chain homotopy. So $N \operatorname{Sym}^3 \Gamma$ turns the chain homotopy between $\operatorname{Tot}(N\Gamma K. \otimes N\Gamma L.)$ and $N\Delta(\Gamma K. \otimes \Gamma L.)$ into a chain homotopy between $N \operatorname{Sym}^3 \Gamma \operatorname{Tot}(N\Gamma K. \otimes N\Gamma L.)$ and $N \operatorname{Sym}^3 \Gamma N\Delta(\Gamma K. \otimes \Gamma L.)$. Chain homotopic complexes are quasi-isomorphic, so continuing our calculation of $G_k(R/I)$ where we left off we get:

$$\begin{aligned} G_k(R/I) &\cong H_k N \operatorname{Sym}^3 \Gamma N\Delta(\Gamma K. \otimes \Gamma L.) \cong H_k N \operatorname{Sym}^3 \Delta(\Gamma K. \otimes \Gamma L.) \\ &\cong H_k N\Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.). \end{aligned}$$

Now we calculate $G_k(R/I)$ by calculating $H_k N\Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.)$. We cannot calculate this directly, so instead we employ the short exact sequences that come from the Cauchy decomposition; these short exact sequences will allow us to get information about the homologies of $N\Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.)$ from easier to calculate homologies.

From the Cauchy decomposition we get the following short exact sequences of bisimplicial modules:

$$0 \rightarrow \Lambda^3 \Gamma K. \otimes \Lambda^3 \Gamma L. \rightarrow M_{(2,1)}(\operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.)) \rightarrow L_1^3 \Gamma K. \otimes L_1^3 \Gamma L. \rightarrow 0$$

$$\begin{aligned} 0 \rightarrow M_{(2,1)}(\operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.)) &\rightarrow \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \\ &\rightarrow \operatorname{Sym}^3 \Gamma K. \otimes \operatorname{Sym}^3 \Gamma L. \rightarrow 0. \end{aligned}$$

Applying $N\Delta$ to this gives us a short exact sequence of chain complexes. We can turn the homologies of these into two long exact sequences, this will allow us to get information about the homologies of $M_{(2,1)}(\operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.))$ from the easier to calculate homologies of $\Lambda^3 \Gamma K. \otimes \Lambda^3 \Gamma L.$ and $L_1^3 \Gamma K. \otimes L_1^3 \Gamma L.$. This information about the homologies of $M_{(2,1)}(\operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.))$ together with the homologies of the easier to calculate homologies of $\operatorname{Sym}^3 \Gamma K. \otimes \operatorname{Sym}^3 \Gamma L.$ will tell us the ranks of the homologies of $\operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.)$.

First we calculate the homologies of $L_1^3 \Gamma K$. The definition of L_1^3 gives us the following short exact sequence for any finitely generated projective module P

$$0 \rightarrow L_1^3 P \rightarrow P \otimes \text{Sym}^2 P \rightarrow \text{Sym}^3 P \rightarrow 0,$$

which gives us the short exact sequence of simplicial complexes

$$0 \rightarrow L_1^3 \Gamma K. \rightarrow \Gamma K. \otimes \text{Sym}^2 \Gamma K. \rightarrow \text{Sym}^3 \Gamma K. \rightarrow 0,$$

the middle term of this short exact sequence is the simplicial complex whose k^{th} term is $\Gamma K_k \otimes \text{Sym}^2 \Gamma K_k$. We think of this middle term instead as the diagonal of a bisimplicial complex whose $(k, l)^{\text{th}}$ term is $\Gamma K_k \otimes \text{Sym}^2 \Gamma K_l$. Now applying the functor N turns this into a short exact sequence of chain complexes

$$0 \rightarrow NL_1^3 \Gamma K. \rightarrow N\Delta(\Gamma K. \otimes \text{Sym}^2 \Gamma K.) \rightarrow N\text{Sym}^3 \Gamma K. \rightarrow 0,$$

and from this we can create a long exact sequence that gives us information about $H_k NL_1^3 \Gamma K.$

Applying the Eilenberg-Zilber Theorem, then Propositions 5.3 and 5.4 we see that:

$$\begin{aligned} H_k N\Delta(\Gamma K. \otimes \text{Sym}^2 \Gamma K.) &\cong H_k \text{Tot}(N\Gamma K. \otimes N\text{Sym}^2 \Gamma K.) \\ &\cong H_k \text{Tot}(K. \otimes \text{Kos}^2(f)). \end{aligned}$$

Now

$$\begin{aligned} \text{Kos}^2(f) &= (\Lambda^2(R) \rightarrow \Lambda^1(R) \otimes \text{Sym}^1(R) \rightarrow \text{Sym}^2(R)) \\ &= (0 \rightarrow R \xrightarrow{f} R) = K., \end{aligned}$$

and therefore

$$\begin{aligned} H_k N\Delta(\Gamma K. \otimes \text{Sym}^2 \Gamma K.) &\cong H_k \text{Tot}(K. \otimes K.) = H_k(P.(R/(f))^{\otimes 2}) \\ &= \Lambda^k((f)/(f)^2) \cong \begin{cases} R/(f) & k = 0, 1 \\ 0 & k > 1 \end{cases} \end{aligned}$$

with the last step given by the isomorphism $H_k(P.(R/J)^{\otimes m}) \cong \Lambda^k((J/J^2)^{m-1})$, see Theorem 5.1 of [Kö].

Using Proposition 5.3 we get

$$H_k N\text{Sym}^3 \Gamma K. \cong H_k(\text{Kos}^3(f))$$

now

$$\begin{aligned} \text{Kos}^3(f) &= (\Lambda^3(R) \rightarrow \Lambda^2(R) \otimes R \rightarrow \Lambda^1(R) \otimes \text{Sym}^2(R) \rightarrow \text{Sym}^3(R)) \\ &= (0 \rightarrow 0 \rightarrow R \xrightarrow{f} R) = K., \end{aligned}$$

hence

$$H_k N \operatorname{Sym}^3 \Gamma K. \cong \begin{cases} R/(f) & k = 0 \\ 0 & k \neq 0. \end{cases}$$

So the long exact sequence of homologies that we get from the short exact sequence

$$0 \rightarrow NL_1^3 \Gamma K. \rightarrow N\Delta(\Gamma K. \otimes \text{Sym}^2 \Gamma K.) \rightarrow N \text{Sym}^3 \Gamma K. \rightarrow 0$$

is

$$\begin{array}{ccccccc}
 & \cdots & H_3NL_1^3\Gamma K. & \longrightarrow & 0 & \longrightarrow & 0 \\
 & & & & & & \downarrow \\
 & & H_2NL_1^3\Gamma K. & \longrightarrow & 0 & \longrightarrow & 0 \\
 & & & & & & \downarrow \\
 & & H_1NL_1^3\Gamma K. & \longrightarrow & R/(f) & \longrightarrow & 0 \\
 & & & & & & \downarrow \\
 & & H_0NL_1^3\Gamma K. & \longrightarrow & R/(f) & \longrightarrow & R/(f) \longrightarrow 0
 \end{array}$$

and so we get

$$H_k N L_1^3 \Gamma K. \cong \begin{cases} 0 & k \neq 1 \\ R/(f) & k = 1. \end{cases}$$

Similarly we get

$$H_k NL_1^3 \Gamma L. \cong \begin{cases} 0 & k \neq 1 \\ R/(g) & k = 1. \end{cases}$$

Now we work with the short exact sequence of simplicial modules

$$0 \rightarrow \Lambda^3 \Gamma K. \otimes \Lambda^3 \Gamma L. \rightarrow M_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \rightarrow L_1^3 \Gamma K. \otimes L_1^3 \Gamma L. \rightarrow 0.$$

As above, we rewrite the left and right hand side of this exact sequence using Δ and we apply the functor N to get the following short exact sequence of chain complexes

$$\begin{aligned} 0 \rightarrow N\Delta(\Lambda^3\Gamma K. \otimes \Lambda^3\Gamma L.) &\rightarrow NM_{(2,1)}(\mathrm{Sym}^3(\Gamma K. \otimes \Gamma L.)) \\ &\rightarrow N\Delta(L_1^3\Gamma K. \otimes L_1^3\Gamma L.) \rightarrow 0. \end{aligned}$$

The Eilenberg-Zilber Theorem tells us that

$$H_k N\Delta(\Lambda^3 \Gamma K. \otimes \Lambda^3 \Gamma L.) \cong H_k \text{Tot}(N\Lambda^3 \Gamma K. \otimes N\Lambda^3 \Gamma L.)$$

and

$$H_k N\Delta(L_1^3 \Gamma K. \otimes L_1^3 \Gamma L.) \cong H_k \text{Tot}(NL_1^3 \Gamma K. \otimes NL_1^3 \Gamma L.).$$

Now by Propositions 5.3 and 5.4 we get

$$H_k \text{Tot}(N\Lambda^3 \Gamma K. \otimes N\Lambda^3 \Gamma L.) \cong H_k \text{Tot}(\tilde{\text{Kos}}^3(f) \otimes \tilde{\text{Kos}}^3(g)).$$

Now

$$\begin{aligned} \tilde{\text{Kos}}^3(f) &= (D^3(R) \rightarrow D^2(R) \otimes R \rightarrow R \otimes \Lambda^2(R) \rightarrow \Lambda^3(R)) \\ &= (R \xrightarrow{f} R \rightarrow 0 \rightarrow 0) = K.[-2], \end{aligned}$$

and similarly $\tilde{\text{Kos}}^3(g) = L.[-2]$. So

$$\begin{aligned} H_k \text{Tot}(N\Lambda^3 \Gamma K. \otimes N\Lambda^3 \Gamma L.) &\cong H_k \text{Tot}(K.[-2] \otimes L.[-2]) \\ &\cong H_k(\text{Tot}(K. \otimes L.)[-4]) \\ &\cong H_{k-4}(P.(R/I)) \cong \begin{cases} 0 & k \neq 4 \\ R/I & k = 4. \end{cases} \end{aligned}$$

Now to calculate $H_k \text{Tot}(NL_1^3 \Gamma K. \otimes NL_1^3 \Gamma L.)$ we note that it is the hyper-tor functor $\mathbf{Tor}_k(NL_1^3 \Gamma K., NL_1^3 \Gamma L.)$ and use the hypertor spectral sequence of Proposition 5.4 taking $A. = NL_1^3 \Gamma K.$ and $B. = NL_1^3 \Gamma L.$. But since $H_k NL_1^3 \Gamma K.$ and $H_k NL_1^3 \Gamma L.$ are 0 unless $k = 1$ (see above) this spectral sequence collapses, with the only (potentially) non-zero terms being when $q' = q'' = 1$, i.e. when $q = 2$. These (potentially) non-zero terms are $\text{Tor}_p(R/(f), R/(g))$. Taking $K.$ as a projective resolution of $R/(f)$ then tensoring throughout by $R/(g)$ we get the chain complex

$$(0 \rightarrow R \otimes R/(g) \xrightarrow{f} R \otimes R/(g)) = (0 \rightarrow R/(g) \xrightarrow{f} R/(g))$$

which has homology R/I at the 0^{th} place and 0 everywhere else. And so

$$H_k(\text{Tot}(NL_1^3 \Gamma K. \otimes NL_1^3 \Gamma L.)) \cong \text{Tor}_{k-2}(R/(f), R/(g)) \cong \begin{cases} 0 & k \neq 2 \\ R/I & k = 2. \end{cases}$$

So the short exact sequence of chain complexes

$$\begin{aligned} 0 \rightarrow N\Delta(\Lambda^3 \Gamma K. \otimes \Lambda^3 \Gamma L.) &\rightarrow NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \\ &\rightarrow N\Delta(L_1^3 \Gamma K. \otimes L_1^3 \Gamma L.) \rightarrow 0 \end{aligned}$$

gives rise to the following long exact sequence of homologies

$$\begin{array}{ccccccc}
\cdots & \rightarrow & 0 & \longrightarrow & H_5 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) & \longrightarrow & 0 \\
& & & & & & \searrow \\
& & & & & & \rightarrow R/I \longrightarrow H_4 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & \rightarrow 0 \longrightarrow H_3 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & \rightarrow 0 \longrightarrow H_2 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \longrightarrow R/I \\
& & & & & & \searrow \\
& & & & & & \rightarrow 0 \longrightarrow H_1 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & \rightarrow 0 \longrightarrow H_0 NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \longrightarrow 0 \longrightarrow 0
\end{array}$$

and therefore we get

$$H_k NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) \cong \begin{cases} R/I & k = 2, 4 \\ 0 & \text{otherwise.} \end{cases}$$

Now we work with the short exact sequence of simplicial modules

$$\begin{aligned}
0 \rightarrow M_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) &\rightarrow \text{Sym}^3(\Gamma K. \otimes \Gamma L.) \\
&\rightarrow \text{Sym}^3 \Gamma K. \otimes \text{Sym}^3 \Gamma L. \rightarrow 0
\end{aligned}$$

the term $\text{Sym}^3 \Gamma K. \otimes \text{Sym}^3 \Gamma L.$ is a simplicial module whose k^{th} place is $\text{Sym}^3 \Gamma K_k \otimes \text{Sym}^3 \Gamma L_k$, but as above it is more useful to think of it as the diagonal of the bisimplicial complex whose $(k, l)^{\text{th}}$ place is $\text{Sym}^3 \Gamma K_k \otimes \text{Sym}^3 \Gamma L_l$. Applying the functor N we get the following short exact sequence of chain complexes

$$\begin{aligned}
0 \rightarrow NM_{(2,1)}(\text{Sym}^3(\Gamma K. \otimes \Gamma L.)) &\rightarrow N \text{Sym}^3(\Gamma K. \otimes \Gamma L.) \\
&\rightarrow N \Delta(\text{Sym}^3 \Gamma K. \otimes \text{Sym}^3 \Gamma L.) \rightarrow 0.
\end{aligned}$$

Applying the Eilenberg-Zilber Theorem and Propositions 5.3 and 5.4 we see

$$\begin{aligned}
H_k N \Delta(\text{Sym}^3 \Gamma K. \otimes \text{Sym}^3 \Gamma L.) &\cong H_k \text{Tot}(N \text{Sym}^3 \Gamma K. \otimes N \text{Sym}^3 \Gamma L.) \\
&\cong H_k \text{Tot}(\text{Kos}^3(f) \otimes \text{Kos}^3(g)).
\end{aligned}$$

Earlier in this proof we showed that $\text{Kos}^3(f) = K$. and similarly $\text{Kos}^3(g) = L$, so

$$H_k \operatorname{Tot}(\operatorname{Kos}^3(f) \otimes \operatorname{Kos}^3(g)) \cong H_k(\operatorname{Tot}(K. \otimes L.)) = H_k(P.(R/I)).$$

Hence

$$H_k N\Delta(\mathrm{Sym}^3 \Gamma K. \otimes \mathrm{Sym}^3 \Gamma L.) \cong \begin{cases} R/I & k = 0 \\ 0 & k \neq 0. \end{cases}$$

And so the short exact sequence of chain complexes

$$\begin{aligned} 0 \rightarrow NM_{(2,1)}(\mathrm{Sym}^3(\Gamma K. \otimes \Gamma L.)) &\rightarrow N \mathrm{Sym}^3(\Gamma K. \otimes \Gamma L.) \\ &\rightarrow N\Delta(\mathrm{Sym}^3(\Gamma K.) \otimes \mathrm{Sym}^3(\Gamma L.)) \rightarrow 0, \end{aligned}$$

gives rise to the following long exact sequence

$$\begin{array}{ccccccc}
\cdots & \rightarrow & 0 & \longrightarrow & H_5 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) & \longrightarrow & 0 \\
& & & & & & \searrow \\
& & & & & & R/I \longrightarrow H_4 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & 0 \longrightarrow H_3 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & R/I \longrightarrow H_2 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & 0 \longrightarrow H_1 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \longrightarrow 0 \\
& & & & & & \searrow \\
& & & & & & 0 \longrightarrow H_0 N \Delta \operatorname{Sym}^3(\Gamma K. \otimes \Gamma L.) \longrightarrow R/I \longrightarrow 0.
\end{array}$$

And hence (as we know $G_k(R/I) \cong H_k N \Delta \text{Sym}^3(\Gamma K. \otimes \Gamma L.)$) we see that

$$G_k(R/I) \cong \begin{cases} R/I & k = 0, 2, 4 \\ 0 & \text{otherwise,} \end{cases}$$

as desired.

☐

References

- [ABW] K. AKIN, D. A. BUCHSBAUM, and J. WEYMAN, *Schur functors and Schur complexes*, Adv. Math. 44 (1982), 207-278.
- [DP] A. DOLD and D. PUPPE, *Homologie nicht-additiver Funktoren. Anwendungen*, Ann. Inst. Fourier Grenoble 11 (1961), 201-312.
- [EM] S. EILENBERG and S. MAC LANE, *On the groups $H(\Pi, n)$, II*, Ann. of Math. 60 (1954), 49-139.
- [JM] B. JOHNSON and R. MCCARTHY, *Linearization, Dold-Puppe stabilization, and Mac Lane's Q -construction*, Trans. Amer. Math. Soc. 350 (1998), 1555-1593.
- [Kö] B. KÖCK, *Computing the homology of Koszul complexes*, Trans. Amer. Math. Soc. 353 (2001), 3115-3147.
- [Ku] E. KUNZ, *Introduction to Commutative Algebra and Algebraic Geometry*, Birkhäuser, Boston (1985).
- [L] S. LANG, *Algebra*, Addison-Wesley, Reading (1971).
- [M] J. P. MAY, *Simplicial Objects in Algebraic Topology*, The University of Chicago Press, Chicago (1967).
- [W] C. A. WEIBEL, *An introduction to homological algebra*, Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge (1994).